



PSYCHALIVE

Welcome to the Webinar, Suicide: What Every Therapist Needs to Know

Lisa Firestone, Ph.D.

Check in through PsychAlive's social media:



Get the latest
news on
PsychAlive
Webinars and
Events



Follow us on Twitter

<http://twitter.com/@psychalive>

Tweet about this webinar: #psychalivewebinar



Like us on Facebook

<http://www.facebook.com/PsychAlive>



THE GLENDON
ASSOCIATION

Suicide: What Every Therapist Needs to Know



with Lisa Firestone, Ph.D.



Lisa Firestone, Ph.D.

Director of Research and Education
The Glendon Association

Senior Editor
PsychAlive.org



THE GLENDON
ASSOCIATION

(For Professionals)
www.glendon.org

PSYCHALIVE

(For the Public)
www.psychalive.org

Glendon@glendon.org



World Suicide Prevention Day

September 10, 2011

International Events – International Association for Suicide Prevention

<http://www.iasp.info/wspd/index.php>

National Suicide Prevention Week

September 4 – 10, 2011

National Events – American Association of Suicidology

<http://www.suicidology.org/web/guest/about-aas/nspw>




Facts About Suicide

- According to the World Health Organization, every 40 seconds a life is lost to suicide, which means that each year we lose nearly 1 million people to suicide.
- For every one person who dies by suicide, 20 or more attempt to end their lives.
- Worldwide, more people die by suicide than from all homicides and wars combined.
- Each person who dies by suicide leaves behind an average of five closely impacted survivors.
- According to a 2008 statistic from SAMHSA, 8.3 million adults in the U.S. had serious thoughts of committing suicide in the past year
- At least 16 percent of American suicides are related to economic strain.
- In 2010 for the second year in a row, more American soldiers, enlisted men, women, and veterans, committed suicide than were killed in wars in Iraq and Afghanistan. The U.S. Army suffered a record number of suicides in July 2011.




Suicidal Thoughts and Behaviors in the Past Year among Adults, by Age Group: 2008



Source: 2008 SAMHSA National Survey on Drug Use and Health (NSDUH).



Suicidal Thoughts and Behaviors in the Past Year among Adults, by Past Year Substance Use Disorder: 2008




Source: 2008 SAMHSA National Survey on Drug Use and Health (NSDUH).




ED Treatment of Mental Disorders

One in 10 suicides are by people seen in the ED within two months of dying.



Mental Health Services

- 19 % of suicides had contact with MH within the past month; 32% within the past year (Luoma, 2002)
- 41% of suicide decedents who had received inpatient psychiatric care died within one year of their discharge; 9% within one day (Pirkis, 1998)
- 71% of psychotherapists report having at least one client who has attempted suicide, while 28% report having had at least one client die by suicide
- Of patients admitted for attempt (Owens et al., 2002)
 - 16% repeat attempts within one year
 - 7% die by suicide within 10 years
 - Risk of suicide “hundreds of times higher” than general population



Poll #1

Have you been impacted by
a suicide attempt or the loss of someone to suicide
either personally or professionally?

Professionally

Personally

Both

Neither



Misconceptions About Suicide

- Most suicides are caused by one particular trigger event.
- Most suicides occur with little or no warning.
- It is best to avoid the topic of suicide.
- People who talk about suicide don't do it.
- Nonfatal self-destructive acts (suicide attempts) are only attention-getting behaviors.




Misconceptions About Suicide

- A suicidal person clearly wants to die.
- Once a person attempts suicide, he or she won't try it again.
- Suicide is a complex problem.
- If a person who has been depressed is suddenly feeling better, the danger of suicide is gone.
- Poor people are the source of most suicides.
- Being religious protects against suicide.



Voices of Suicide

From "Voices of Suicide: Learning From Those Who Lived"



http://www.youtube.com/watch?v=Es7s_z-YVLE



Poll #2

Have you had clients who've reported these types of thoughts?

- *You are so stupid. You never get anything right.*
- *You are different from other people.*
- *You're so unattractive.*




Our Approach to Suicide

Each person is divided:

- One part wants to live and is goal directed and life affirming.
- And one part is self-critical, self-hating and at its ultimate end, self-destructive. The nature and degree of this division varies for each individual.

Real Self - Positive



Anti-Self - Critical





Division of the Mind

Self-System

Positive Parenting Behaviors
Attunement, Affection, Control
Parental Nurturance/Genetic
Predisposition/Temperament

Other nurturing experiences:
Caring adults, education



Anti-Self System

Destructive Parental Behaviors/ Misattunement,
lack of affection, reject, neglect, hostility,
permissiveness, Genetic Predisposition,
Temperament

Other factors: accidents, illnesses, traumatic separation





Self System

Unique make-up of the individual –harmonious identification and incorporation of parent's positive attitudes and traits and the effect of experience on the maturing self-system resulting in a stronger sense of self and a greater degree of differentiation from parents or early caretakers.





Personal Attitudes/Goals/Conscience

Realistic, Positive Attitudes Towards Self

Realistic evaluation of talents, abilities, etc...with generally positive/ compassionate attitude towards self and others.

Goals

Needs, wants, search for meaning in life

Moral Principles



Behavior

Ethical behavior towards self and others

Goal Directed Behavior

Acting with Integrity





The Anti-Self System

The Fantasy Bond (core defense) furthers a self-parenting process made up of both the helpless, needy child, and the self-punishing, self-nurturing parent. Either aspect may be acted out in romantic relationships. The degree of reliance on this defense is proportional to the amount of damage sustained while growing up.






The Self-Parenting Process Self-Punishing Voices

Voice Process

Critical thoughts toward self



2. Micro-suicidal injunctions
(punitive/cruel thoughts)



3. Suicidal injunctions
– suicidal ideation




Behaviors

Verbal attacks-a generally negative
attitude toward self and others
predisposing alienation



Addictive Patterns/thoughts
heightening emotional distress

Actions that jeopardize one's
health and safety; physical attacks
on the self, and actual suicide



The Self-Parenting Process Self-Soothing Voices

Voice Process



1. Self Soothing Attitudes
2. Micro-suicidal injunctions (seductive/self-indulgent thoughts)
3. Aggrandizing thoughts towards self
4. Suspicious paranoid thoughts toward others

Behaviors

Self-limiting or self-protective lifestyles, Inwardness

Addictive patterns/ numbing emotional distress


Narcissism/ Vanity and aggressive behavior towards others

Alienation, hostile attitudes towards others and aggressive behavior




Developmental

From "Voices of Suicide: Learning From Those Who Lived"




<http://www.youtube.com/watch?v=xPcgkm9AlhU>



Our Approach to Suicide


Negative thoughts exist on a continuum, from mild self-critical thoughts to extreme self-hatred to thoughts about suicide






Our Approach to Suicide

Self-destructive behaviors exist on a continuum from self-denial to substance abuse to actual suicide.






Our Approach to Suicide

There is a relationship between these two continuums. How a person is thinking is predictive of how he or she is likely to behave.





Definition of the Voice

The critical inner voice refers to a well-integrated pattern of destructive thoughts toward our selves and others. The “voices” that make up this internalized dialogue are at the root of much of our maladaptive behavior. This internal enemy fosters inwardness, distrust, self-criticism, self-denial, addictions and a retreat from goal-directed activities. The critical inner voice effects every aspect of our lives: our self-esteem and confidence, our personal and intimate relationships, and our performance and accomplishments at school and work.






Continuum of Negative Thought Patterns

Thoughts that lead to low-self-esteem or inwardness (self-defeating thoughts):

Levels of Increasing Suicidal Intention

Content of Voice Statements

- | | | |
|---|---|--|
| 1. Self-deprecating thoughts of everyday life | ↔ | <i>You're incompetent, stupid. You're not very attractive. You're going to make a fool of yourself.</i> |
| 2. Thoughts rationalizing self-denial; thoughts discouraging the person from engaging in pleasurable activities | ↔ | <i>You're too young (old) and inexperienced to apply for this job. You're too shy to make any new friends. Why go on this trip? It'll be such a hassle. You'll save money by staying home.</i> |
| 3 Cynical attitudes towards others, leading to alienation and distancing | ↔ | <i>Why go out with her/him? She's cold, unreliable; she'll reject you. She wouldn't go out with you anyway. You can't trust men/women.</i> |



Continuum of Negative Thought Patterns

Thoughts that lead to low-self-esteem or inwardness (self-defeating thoughts):

Levels of Increasing Suicidal Intention

Content of Voice Statements

4. Thoughts influencing isolation; rationalizations for time alone, but using time to become more negative toward oneself




Just be by yourself. You're miserable company anyway; who'd want to be with you? Just stay in the background, out of view.

5. Self-contempt; vicious self-abusive thoughts and accusations (accompanied by intense angry affect)



You idiot! You bitch! You creep! You stupid shit! You don't deserve anything; you're worthless.



Continuum of Negative Thought Patterns

Thoughts that support the cycle of addiction (addictions):


Levels of Increasing Suicidal Intention



Content of Voice Statements

6. Thoughts urging use of substances or food followed by self-criticisms (weakens inhibitions against self-destructive actions, while increasing guilt and self-recrimination following acting out).

*It's okay to do drugs, you'll be more relaxed. Go ahead and have a drink, you deserve it.
(Later) You weak-willed jerk!
You're nothing but a drugged-out drunken freak.*



Continuum of Negative Thought Patterns

Thoughts that lead to suicide (self-annihilating thoughts):

Levels of Increasing Suicidal Intention

- 7. Thoughts contributing to a sense of hopelessness urging withdrawal or removal of oneself completely from the lives of people closest.**
- 8. Thoughts influencing a person to give up priorities and favored activities (points of identity).**
- 9. Injunctions to inflict self-harm at an action level; intense rage against self.**

Content of Voice Statements

See how bad you make your family (friends) feel. They'd be better off without you. It's the only decent thing to do; just stay away and stop bothering them.

What's the use? Your work doesn't matter any more. Why bother even trying? Nothing matters anyway.

Why don't you just drive across the center divider? Just shove your hand under that power saw!



Continuum of Negative Thought Patterns

- Thoughts that lead to suicide (self-annihilating thoughts):

- Levels of Increasing Suicidal Intention

- 10. Thoughts planning details of suicide (calm, rational, often obsessive, indicating complete loss of feeling for the self).
- 11. Injunctions to carry out suicide plans; thoughts baiting the person to commit suicide (extreme thought constriction).

- Content of Voice Statements


- *You have to get hold of some pills, then go to a hotel, etc.*
- *You've thought about this long enough. Just get it over with. It's the only way out.*






Poll #3


Have you had clients express these types of self-destructive thoughts?



Those Who Desire Suicide




Joiner, Thomas. *Why People Die By Suicide*. "The Three Components of Completed Suicide." Harvard University Press, 2005.



Thoughts to Actions

From "Voices of Suicide: Learning From Those Who Lived"



<http://www.youtube.com/watch?v=IMR1I2Gu5Vo>



Suicide Warning Signs

- Disturbed sleep patterns
- Anxiety, agitation
- Pulling away from friends and family
- Past attempts
- Extremely self-hating thoughts
- Feeling like they don't belong
- Hopelessness
- Rage





Suicide Warning Signs

- Feeling trapped
- Increased use of alcohol or drugs
- Feeling that they are a burden to others
- Loss of interest in favorite activities -“nothing matters”
- Giving up on themselves
- Risk-taking behavior
- Suicidal thoughts, plans, actions
- Sudden mood changes for the better





Poll #4

Do you use objective measures to
assess suicide risk?

Why Use Objective Measures?

What Interferes with Clinical Judgment

- Anxiety
- Counter Transference
- Psych Ache
- Research Minimizing
- Diverse Menu of Risk Factors

Our Measures

Based on Separation Theory developed by Robert W. Firestone, PhD. and represents a broadly based coherent system of concepts and hypothesis that integrates psychoanalytic and existential systems of thought. The theoretical approach focuses on **internal negative thought processes**. These thoughts (i.e. “voices”) actually direct behavior and, thus, are likely to predict how an individual will behave.



Firestone Assessment of Self-Destructive Thoughts

	Never	Rarely	Once In A While	Frequently	Most Of The Time
1. Just stay in the background.	0	1	2	3	4
2. Get them to leave you alone. You don't need them.	0	1	2	3	4
3. You'll save money by staying home. Why do you need to go out anyway?	0	1	2	3	4
4. You better take something so you can relax with those people tonight.	0	1	2	3	4
5. Don't buy that new outfit. Look at all the money you are saving.	0	1	2	3	4

Figure 4.3. Mean T Scores for the Depression Sample: Inpatients and Outpatients—Ideators Versus Nonideators (N = 296)





Figure 4.5. Mean *T* Scores for the Bipolar Disorder Sample—Ideators Versus Nonideators (*N* = 68)



**Figure 4.7. Mean T Scores for the Substance Abuse Sample:
Inpatients and Outpatients—Ideators Versus Nonideators
(N = 202)**




Figure 4.6. Mean *T* Scores for the Inpatient Schizophrenia Sample—Ideators Versus Nonideators (*N* = 115)






Figure 4.4. Mean T Scores for the Outpatient Personality Disorder Sample—Ideators Versus Nonideators (N = 35)



Uses for Our Measures

- Risk Assessment
- Treatment Planning
- Targeting Intervention
- Outcome Evaluation

Figure 3. Approximate ROC Curves for the FVSSDB, SPS, and BHS





Suicide Risk Factors Checklist

- Psychiatric

- Major Depression-particularly endogenous
- Alcohol dependence-rate 50x the general population, 25% of all suicides
- Drug addiction- 10% die by suicide
- Personality Disorders- especially borderline or compulsive
- Schizophrenia-frequently with command hallucinations
- Organic psychoses



Suicide Risk Factors Checklist

- Psychiatric

- Past history- especially if attempts were serious
- Family History-increased risk in twin and adoption studies
- Possible biologic markers: Decreased CSF 5-HIAA, increased CSF MHPG, nonsuppressing DST, low platelet MAO, low platelet serotonin, high platelet serotonin-2 receptor responsibility
- Poor physical health- renal dialysis patients have a suicide rate 400X higher than the general population



Suicide Risk Factors Checklist

- Psychological
 - History of Recent Loss
 - History of parental Loss During Childhood
 - Important Days-anniversaries, holidays, etc.
 - Family instability
 - Social Isolation-loss of social supports



Suicide Risk Factors Checklist

- Social
 - Sex-
Male 3X female
 - Race-
Whites 2x nonwhites, except urban areas where rate is the same: Native Americans have higher rates
 - Age-
in men rates rise with age above age 45; in women the peak risk is about age 55, then the rate declines
 - Religion-
Protestants and atheists have higher rates than Jews and Catholics



Suicide Risk Factors Checklist

- Social
 - Geography-
urban rates higher
 - Marital Status-
divorced> single>widowed>married
 - Socioeconomic-
high rates at both spectrums, retired and
unemployed at higher risk



Examination of a Potentially Suicidal Patient

- Is there a wish to die?
- Is there a plan?
- What is the method planned?
- What epidemiologic risk factors are present?
- Is there a history of recent substance use?
- What medical illnesses are present?
- What psychiatric diagnoses are present?



Examination of a Potentially Suicidal Patient

- Is there a past or family history of suicide attempts?
- Is there a history of impulsivity.
- What is the level of psychological defensive functioning?
- Has there been a will made recently?
- Is there a history of recent losses, and how do they relate to past history of losses?
- Is there talk of plans for the future?
- What is the nature of the patient's social support system?



Protective Factors

- Family and community connections/ support
- Clinical Care (availability and accessibility)
- Resilience
- Coping Skills
- Frustration tolerance and emotion regulation
- Cultural and religious beliefs; spirituality





Poll #5

Do you find it stressful to assess
and manage suicide risk?

- Yes, very
- Yes, somewhat
- No




Figure 5.1. Assessment and Management of the Suicidal Patient




Figure 5.1. Assessment and Management of the Suicidal Patient

CAMS

SUICIDE STATUS FORM-III (SSF III) INITIAL SESSION

Patient: _____ Clinician: _____ Date: _____ Time: _____

Section A (Patient):

Rate and fill out each item according to how you feel right now.

Rank Then rank items in order of importance 1 to 5 (1 = most important to 5 = least important).

_____	1) RATE PSYCHOLOGICAL PAIN (<i>hurt, anguish, or misery in your mind; <u>not</u> stress; <u>not</u> physical pain</i>): Low Pain: 1 2 3 4 5 :High Pain What I find most painful is: _____
_____	2) RATE STRESS (<i>your general feeling of being pressured or overwhelmed</i>): Low Stress: 1 2 3 4 5 :High Stress What I find most stressful is: _____
_____	3) RATE AGITATION (<i>emotional urgency; feeling that you need to take action; <u>not</u> irritation; <u>not</u> annoyance</i>): Low Agitation: 1 2 3 4 5 :High Agitation I most need to take action when: _____
_____	4) RATE HOPELESSNESS (<i>your expectation that things will not get better no matter what you do</i>): Low Hopelessness: 1 2 3 4 5 :High Hopelessness I am most hopeless about: _____
_____	5) RATE SELF-HATE (<i>your general feeling of disliking yourself; having no self-esteem; having no self-respect</i>): Low Self-Hate: 1 2 3 4 5 :High Self-Hate What I hate most about myself is: _____
N/A	6) RATE OVERALL RISK OF SUICIDE: Extremely Low Risk: 1 2 3 4 5 :Extremely High Risk (will <u>not</u> kill self) (will kill self)

- 1) How much is being suicidal related to thoughts and feelings about yourself? **Not at all: 1 2 3 4 5 :completely**
- 2) How much is being suicidal related to thoughts and feelings about others? **Not at all: 1 2 3 4 5 :completely**


Please list your reasons for wanting to live and your reasons for wanting to die. Then rank in order of importance 1 to 5.

Rank	REASONS FOR LIVING	Rank	REASONS FOR DYING

I wish to live to the following extent: **Not at all: 0 1 2 3 4 5 6 7 8 :Very much**

I wish to die to the following extent: **Not at all: 0 1 2 3 4 5 6 7 8 :Very much**

The one thing that would help me no longer feel suicidal would be: _____



Poll #6

Do you see clients who have had prior suicide attempts?

- No past attempts
- 1 past attempt
- 2 or more past attempts



Multiple Attempters as a Special High-Risk Group (in comparison to single attempters/ideators)

- Distinctive in every way
 - Greater likelihood to have diagnosis, co-morbidity, personality disorder
 - Younger at time of first attempt (greater chronicity)
 - Lower lethality first attempt (raises question about intent, function of behavior)
 - More impulsive
 - More likely to be associated with substance abuse
 - Greater symptom severity
 - Anxiety, depression, hopelessness, anger, suicidal ideation (frequency, intensity, specificity, duration, intent)
 - More frequent histories of trauma, abuse
 - Distinctive characteristics of crises




Figure 3.3. Suicide-relevant cognitions and the likelihood of engaging in suicidal acts.




SAFETY PLAN	
1. Warning signs (when I am to use the safety plan): <input type="checkbox"/> Wanting to go to sleep and not wake up <input type="checkbox"/> Wanting to hurt myself <input type="checkbox"/> Thinking "I can't take it anymore"	
2. Coping strategies (things I can try to do on my own): <input type="checkbox"/> Listening to rock music <input type="checkbox"/> Rocking in chair <input type="checkbox"/> Going for a walk <input type="checkbox"/> Controlled breathing <input type="checkbox"/> Taking a hot or cold shower <input type="checkbox"/> Exercising	
3. Contacting other people: <input type="checkbox"/> Calling a friend to distract myself: _____ Phone: _____ If distraction does not work, I will tell any of the following people that I am in crisis and ask for help: <input type="checkbox"/> Calling a family member: _____ Phone: _____ <input type="checkbox"/> Calling or talking to someone else: _____ Phone: _____	
4. Contacting a health care professional during business hours: <input type="checkbox"/> Calling my therapist: _____ Phone: _____ <input type="checkbox"/> Calling my psychiatrist: _____ Phone: _____ <input type="checkbox"/> Calling my case manager: _____ Phone: _____ The following agencies or services may be called 24 hours a day/7 days a week: <input type="checkbox"/> Calling the psychiatric ED: _____ Phone: _____ <input type="checkbox"/> Calling National Suicide Prevention Lifeline Phone: 1-800-273-TALK	
Patient Signature: _____ Date: _____	
Clinician Signature: _____ Date: _____	

Figure 6.3. Example of a safety plan developed during the early phase of treatment. ED = emergency department.

Safety Plan

Stanley and Brown, 2008



Commitment to Treatment Statement in Practice

We recommend that the CTS always be handwritten and individualized by the clinician; avoid using a standard preprinted form. The CTS should always include a crisis response plan, that is, the specific steps the patient should take during a crisis. Some manner of agreement should be accomplished in the first session. The implicit, and potentially problematic, messages are likely profound with use of a preprinted form. In addition to the central elements noted, it is important to identify any time restrictions imposed by the patient: What is the duration of the agreement – 1 week, 1 month, 1 year? Here is an example of a CTS from our practice (Rudd, Joiner, & Rajab, 2004):

I, _____, agree to make a commitment to the treatment process. I understand that this means that I have agreed to be actively involved in all aspects of treatment including:



10 Most Common Errors in Suicide Prevention

1. Superficial Reassurance
2. Avoidance of Strong Feelings
3. Professionalism
4. Inadequate Assessment of suicidal intent
5. Failure to identify the precipitating event
6. Passivity
7. Insufficient Directiveness
8. Advice Giving
9. Stereotypic Responses
10. Defensiveness



Practice Recommendations

1. When imminent risk does not dictate hospitalization, the intensity of outpatient treatment (i.e., more frequent appointments, telephone contacts, concurrent individual and group treatment) should vary in accordance with risk indicators for those identified as high risk.




Practice Recommendations

2. If the target goal is a reduction in suicide attempts and related behaviors, treatment should be conceptualized as long-term and target identified skills deficits (e.g., emotion regulation, distress tolerance, impulsivity, problem-solving, interpersonal assertiveness, anger management), in addition to other salient treatment issues.



Practice Recommendations

3. If therapy is brief and the target variable are suicidal ideation, or related symptomatology such as depression, hopelessness, or loneliness, a problem-solving component should be used in some form or fashion as a core intervention.




Practice Recommendations

4. Regardless of therapeutic orientation, an explanatory model should be detailed identifying treatment targets, both direct (i.e., suicidal ideation, attempts, related self-destructive and self-mutilatory behaviors) and indirect (depression, hopelessness, anxiety, and anger; interpersonal relationship dysfunction; low self-esteem and poor self-image; day-to-day functioning at work and home).



Practice Recommendations


5. Use of standardized follow-up and referral procedure (e.g., letters or phone calls) is recommended for those dropping out of treatment prematurely in an effort to enhance compliance and reduce risk for subsequent attempts.



Poll #7

Have you received formal training in treating suicidal individuals?

- None
- Some
- A comprehensive course



Effective Therapy Approaches for Treating the Suicidal Person

- Cognitive Therapy for suicidal people was developed by Aaron Beck and Gregory Brown. Unlike other CBT treatments, this approach is not time limited. The third and last stage is "Relapse Prevention with a Twist." Clients do not graduate from treatment until they demonstrate that they are ready to do this on their own.
- Dialectical Behavior Therapy, developed by Marsha M. Linehan, is designed to treat emotion regulation difficulties and suicidal behavior. One element, the skill-building component of DBT, addresses the issues of distress tolerance and the development of healthy affect regulation strategies, both of which are essential for suicidal clients.
- Mentalizing Treatment, developed by Jon Allen and Peter Fonagy, emphasizes emotional regulation and expressiveness. The techniques implemented assist clients in forming good affect regulation and tolerance through the process of developing the mentalizing capability to observe and understand their mind and the minds of others, accurately seeing the mind behind the behavior.
- Transference Focused Therapy, developed by Kernberg, Clarkin, and Yeomans, concentrates on the intermediate interaction between the client and therapist in session by focusing on the therapeutic relationship.
- Voice Therapy, which was developed by Robert Firestone, is a cognitive-affective-behavioral therapeutic methodology that brings introjected hostile thoughts, with the accompanying negative affect, to consciousness, rendering them accessible for treatment. This technique facilitates the identification of the negative cognitions driving the suicidal actions, which in turn helps clients to gain a measure of control over all aspects of their self-destructive or suicidal behavior. This process helps clients expand their personal boundaries, develop a sense of meaning in life, and reduce the risk of self-destructive behavior, including suicide.



Construction of a Hope Kit*

Another activity that is undertaken in the middle phase of therapy is the construction of a hope kit. A hope kit consists of a container that holds mementos (photographs, letters, souvenirs) that serve as reminders of reasons to live. Patients are instructed to be as creative as possible when creating their hope kit, so that the end result is a powerful and personal reminder of their connection to life that can be used when feeling suicidal. We have found that patients report making their hope kits to be a highly rewarding experience that often leads them to discover reasons to live they had previously overlooked.



Suzanne was rather artistic and reported that she enjoyed this task. She found an old shoe box and decorated it using some of her favorite pictures. Inside she included pictures of her mother, her friends, and her cat. She also included the lyrics of her favorite song, a potpourri bag filled with her favorite scent, and a piece of her childhood blanket. Suzanne kept the hope box on her dresser, and it frequently reminded her of all the good things in her life.

**Excerpted from "Cognitive Therapy, Cognition, and Suicidal Behavior" by GK Brown, E Jeglic, GR Henriques, and AT Beck In T.E. Ellis (Ed.), *Cognition and Suicide* (APA Books, 2006).



Voice Therapy

Cognitive/Affective/Behavioral Approach





• Self



• Anti-Self






• Interpersonal Neurobiology

- C urious
- O pen
- A ccepting
- L oving



Treatment

From "Voices of Suicide: Learning From Those Who Lived"




<http://www.youtube.com/watch?v=H4i9G9g-1kk>



Most Helpful Aspects from Client Perspective

Validating Relationships


- Participants describe the existence of an affirming and validating relationship as a catalyst for reconnection with others and with oneself. A difficult part of the recovery process was breaking through, cognitive, emotional, and behavioral barriers that participants had generated for survival.



Most Helpful Aspects from Client Perspective

Working with Emotions

- Dealing with the intense emotions underlying suicidal behavior was perceived as crucial to participant's healing. The resolution of despair and helplessness was a pivotal and highly potent experience for all participants in the study. Almost paradoxically, if a client did not receive acknowledgement of these powerful and overwhelming feelings, they reported being unable to move beyond them.



Most Helpful Aspects from Client Perspective

Developing Autonomy and Identity

- Participants identified understanding suicidal behaviors , developing self-awareness, and constructing personal identity as key components of the therapeutic process. Participants conceptualized the therapeutic experience as confronting and discarding negative patterns while establishing new, more positive ones.



Poll #8

Have you received training on the **Standard of Care** for treating suicidal clients?



Practice and Risk Management Guidelines

For each patient seen there must be:

- an initial evaluation and assessment
- regular ongoing clinical evaluations and case reviews
- consultation reports and supervision reports
- formal treatment plan
- a formal informed consent for treatment
- formal assessment of competence
- documentation of confidentiality considerations



Practice and Risk Management Guidelines

1. Documentation

- “If it isn’t written down, it didn’t happen”
- a knowledge of the epidemiology, risk factors, and treatment literature for the suicidal patient

2. Information on previous treatment

- All previous treatment records

3. Involvement of the family and significant others



Practice and Risk Management Guidelines

4. Consultation on present clinical circumstances

- Routinely obtain consultation and/or supervision (or make referrals)
- The principle that two perspectives are better than one should always guide the clinician in moments of clinical uncertainty.

5. Sensitivity to medical issues.

6. Knowledge of community resources

7. Consideration of the effect on self and others



Self-Care & Help Seeking Behaviors

- Ask for help
- Talk to others
- Get plenty of rest
- Drink plenty of water, avoid caffeine
- Do not use alcohol and other drugs
- Exercise
- Use relaxation skills



American Association of Suicidology's Survivors' Support Group Directory

<http://www.suicidology.org/web/guest/support-group-directory>



IASP Suicide Survivor Organizations (listed by country) -

http://www.iasp.info/resources/Postvention/National_Suicide_Survivor_Organizations/

National Suicide Prevention Lifeline - 1-800-273-TALK(8255)

<http://www.suicidepreventionlifeline.org/>




Guidelines for Suicide Counseling


- Directly Explore Suicidal Thoughts
- Assess suicide risk
- Attempt to diffuse the potential for the lethality
- Encourage social connection
- Implement Behavioral Contracting
- Seek Consultative Support
- Make Appropriate Referrals
- Take Suicidal Gestures Seriously
- Remember the healing power of therapeutic availability



Resources: Books



For Public and Professionals



For Professionals


Visit www.psychalive.org for resource links




Resources: Films



For the Public



For Professionals



For Survivors

Visit www.psychalive.org for resource links



Online CE Course



Professional Psych Seminars
Suicide: What Therapists Need to Know
6 CE Units

[http://www.psychsem.com/live/suicide--
what-therapists-need-to-know.php](http://www.psychsem.com/live/suicide--what-therapists-need-to-know.php)



Upcoming Webinars on Suicide Prevention



Understanding and Preventing Suicide

Tuesday, September 27, 2011

11:00 AM - 12:00 PM PDT

Learn more or register here

http://www.psychalive.org/2011/04/psychalive_webinars/

Visit www.psychalive.org for more information or to register for additional webinars.



Upcoming Webinars Featuring Dr. Pat Love



Love in the Time of Twitter: Keeping Relationships Strong in the Age of Social Media

Tuesday, September 20, 2011

11:00 AM - 12:00 PM PDT

Learn more or register here -

http://www.psychalive.org/2011/04/psychalive_webinars/

Visit www.psychalive.org for more information or to register for additional webinars.



Upcoming Webinars Featuring Dr. Donald Meichenbaum



Road Map to Resilience : Ways to Bolster
Resilience and Well-being

Monday, November 14, 2011

11:00 AM - 12:00 PM PDT

Learn more or register here

http://www.psychalive.org/2011/04/psychalive_webinars/



Treatment of Individuals with PTSD, Complex PTSD
and Comorbid Disorders: A Life-Span Approach (CE)

Tuesday, November 15, 2011

11:00 AM - 12:30 PM PDT

Learn more or register here -

http://www.psychalive.org/2011/04/psychalive_webinars/

Visit www.psychalive.org for more information or to register for additional webinars.



Upcoming Webinars on Overcoming Depression



Innovative Approach to Treating Depression (CE)

Tuesday, October 11, 2011

11:00 AM - 12:30 PM PDT

Learn more or register here

http://www.psychalive.org/2011/04/psychalive_webinars/



The Critical Inner Voice That Causes Depression

Tuesday, December 6, 2011

11:00 AM - 12:00 PM PDT

Learn more or register here

http://www.psychalive.org/2011/04/psychalive_webinars/

Visit www.psychalive.org for more information or to register for additional webinars.



Help Support Glendon

Make a donation at



THE GLENDON
ASSOCIATION

ERROR: undefined
OFFENDING COMMAND: f'~

STACK: